

Regimen schedule

Next course of treatment, day 15														
Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Metotrexat	X		X		X		X							
Kalciumfolinat (vattenfritt)		X		X		X		X						

About your treatment

Methotrexate is a cytostatic drug. Calcium folinate is a folic acid, which reduces the risk of side effects from methotrexate.

Each course is 14 days. Methotrexate is given by injection on days 1, 3, 5 and 7. Calcium folinate is given as tablets on days 2, 4, 6 and 8.

During treatment, your blood tests and any side effects will be monitored.

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips: Try to exercise and move around. It's the only thing known to help with fatigue.

Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

Tips: Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

Tips: Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

Tip: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

Problems with mouth and mucous membranes

The mucous membranes of your nose, mouth, eyes and abdomen may be affected by your treatment. For example, you may have nosebleeds and a runny nose. In the mouth, you may experience dryness, redness, burning and sores. You can also get fungus, which turns the lining of your mouth red or gives it a white coating. Eyes may become dry and watery. The mucous membranes in the lower abdomen can also become dry.

Tip: If you have a dry mouth, use saliva stimulants available in pharmacies. Rinsing your mouth with Vichy water can prevent fungus. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist of your cancer treatment when you visit. Avoid contact lenses if you have eye problems. Women can use oestrogen cream, emollient cream or oil to treat dry mucous membranes in the lower abdomen.

Contact your clinic immediately at:

- fever above 38 degrees, or temperature below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- chest pain or difficulty breathing
- nosebleeds or other bleeding that you can't stop
- skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

Common advice

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes
