

**Azacitidin-Venetoklax kur 1.**  
for Haematological malignancy

**Regimen schedule**

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Venetoklax	X																				
Venetoklax		X																			
Venetoklax			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Azacitidin Suspension	X	X	X	X	X																

**Next course of treatment, day 36**

Day	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Venetoklax														
Venetoklax														
Venetoklax														
Azacitidin Suspension														

**About your treatment**

Venetoclax blocks proteins that regulate the growth of cancer cells, slowing it down.  
Take the tablets once a day without interruption.  
During treatment, your blood tests and any side effects will be monitored.

**How to take the medicines**

Swallow the tablets whole with water at mealtime. If you miss a dose, take it only if there are more than 8 hours left until your next scheduled dose.  
You must not eat grapefruit, bitter orange or star fruit (carambola) during treatment.  
Read more in the package leaflet.

**Side effects - symptoms and tips**

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.  
Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

**Fatigue**

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.  
**Tips:** Try to exercise and move around. It's the only thing known to help with fatigue.

## Cough and shortness of breath

You may experience coughing and shortness of breath. Tell your doctor or nurse if you have trouble breathing.

## Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

**Tip:** Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

## Bleeding

You may have an increased risk of bleeding. For example, you may bleed easily from your nose or mouth. If you get a cut on your skin, you may bleed longer than usual. Surgical wounds may heal more slowly.

**Tips:** Be careful not to cut yourself. You can stop nosebleeds by squeezing the soft parts of your nose for 10 minutes. If this doesn't stop it, contact your doctor. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist or other health care provider of your cancer treatment when you visit.

## Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

**Tips:** Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

## Skin problems

You may experience itching and a rash.

**Tip:** When washing your skin, be gentle and use mild soap. Lubricate with moisturiser.

## Contact your clinic immediately at:

- fever above 38 degrees, or temperature below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- chest pain or difficulty breathing
- nosebleeds or other bleeding that you can't stop
- skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

## Common advice

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## Notes

---

---