

## Palbociklib - Kapsel

for Breastcancer

### Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Palbociklib	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

### Next course of treatment, day 29

Day	22	23	24	25	26	27	28
Palbociklib							

## About your treatment

Palbociclib blocks proteins that regulate the growth and division of cancer cells, slowing it down. Palbociclib (Ibance) is taken as capsules. It is given together with hormone treatments for cancer.

Each course is 28 days. You take capsules once a day from day 1 to 21. After that, there is a 7-day break, before a new course starts. Your doctor or nurse will tell you how many capsules to take at a time. During treatment, your blood tests and any side effects will be monitored.

## How to take the medicines

Swallow the capsules whole with water. Take the capsules with a meal at about the same time each day. If you miss a dose or vomit, take your usual dose at the next opportunity. Grapefruit and grapefruit juice should be avoided as this may increase side effects. Also St. John's wort should be avoided as it affects the effect of the drug

Read more in the package leaflet.

## Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

### Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

**Tips:** Try to exercise and move around. It's the only thing known to help with fatigue.

## Problems with mouth and mucous membranes

The mucous membranes of your nose, mouth, eyes and abdomen may be affected by your treatment. For example, you may have nosebleeds and a runny nose. In the mouth, you may experience dryness, redness, burning and sores. You can also get fungus, which turns the lining of your mouth red or gives it a white coating. Eyes may become dry and watery. The mucous membranes in the lower abdomen can also become dry.

**Tips:** If you have a dry mouth, use saliva stimulants available in pharmacies. Rinsing your mouth with Vichy water can prevent fungus. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist of your cancer treatment when you visit. Avoid contact lenses if you have eye problems. For dry mucous membranes in the lower abdomen, vaginal gel, vagitories and emollient cream (without oestrogen) can be used.

## Nausea and taste changes

You may feel unwell from the treatment. You will therefore be given anti-nausea medication in conjunction with the treatment. The medicines may cause constipation. You can get different varieties depending on how you feel. The taste of food and drink may be altered by the treatment.

**Tips:** Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

## Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

**Tips:** Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

## Eye problems

You may get red eyes with stinging and pain. You may also experience blurred vision, dry eyes and become more sensitive to light.

**Tip:** If your eyes sting and feel dry, use lubricating eye drops available from pharmacies.

## Contact your clinic immediately at

### Acute

Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath

### Contact your clinic immediately at:

- Fever (temperature above 38 degrees)
- Sudden deterioration regardless of symptoms

## Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## Notes

---

---