

## PCV

for Tumour in the brain and spinal cord

## Patient information

### Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Lomustin	X																				
Procarbazine								X	X	X	X	X	X	X	X	X	X	X	X	X	X
Vinkristin								X													

### Next course of treatment, day 43

Day	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42
Lomustin																					
Procarbazine																					
Vinkristin								X													

## About your treatment

Lomustine, procarbazine and vincristine are cytostatic drugs.

Each course is 42 days. On day 1, you take Lomustine capsules. On days 8-21, you take procarbazine capsules. Your doctor or nurse will tell you how many capsules to take at a time. On days 8 and 29, you will be given an IV. After that, there is a break of 13 days, before a new course starts.

During treatment, your blood tests and any side effects will be monitored.

## How to take the medicines

Lomustine: Swallow the capsules whole with water, preferably at bedtime or three hours after a meal.

Procarbazine: Swallow the capsules whole with water at bedtime, with or without food. Foods high in tyramine, such as mouldy cheese, banana and yoghurt, should be avoided. Procarbazine has an antabuse-like effect, so you should not drink alcohol during treatment.

If you vomit after taking a capsule, do not take new capsules the same day.

Read more in the package leaflet.

## Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

### Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

**Tips:** Try to exercise and move around. It's the only thing known to help with fatigue.

## **Nausea and taste changes**

You may feel unwell from the treatment. You will therefore be given anti-nausea medication in conjunction with the treatment. The medicines may cause constipation. You can get different varieties depending on how you feel. The taste of food and drink may be altered by the treatment.

**Tips:** Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

## **Problems with stomach and intestines**

Constipation is common. You may also get diarrhoea.

**Tip:** Drink a lot, a few glasses more than usual per day. If you're constipated, try high-fibre foods and exercise. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

## **Infection susceptibility**

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

**Tips:** Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

## **Problems with skin, hair and nails**

Skin often becomes dry and more sensitive to the sun. You may also lose hair or your hair may change texture. Nails can become brittle.

**Tip:** When washing your skin, be gentle and use mild soap. Lubricate often with moisturiser. Wear covering clothing to protect against the sun, and sunscreen on parts of your body that you can't cover. Wear gloves for work that is dirty or abrasive for your hands. If your nails are bothering you, you can use nail polish and brush your cuticles with oil or apply a moisturiser.

## **Numbness and tingling - sensory disturbances**

You may experience numbness and tingling in your hands and feet, and your fine motor skills may be impaired. For example, you may find it difficult to press buttons or walk. Symptoms usually go away, but not completely in everyone. Tell your doctor or nurse if you have numbness or tingling in your hands and feet.

## **Contact your clinic immediately at:**

- fever above 38 degrees, or temperature below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- chest pain or difficulty breathing
- nosebleeds or other bleeding that you can't stop
- skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

## **Common advice**

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## Notes

---

---