# Brigatinib (underhållsdos)

for Lungcancer

#### Regimen schedule

| Day        | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 |
|------------|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|
| Brigatinib | Х | Х | Х | Х | Х | Х | Х | Х | Х | Х  | Х  | Х  | Χ  | Х  | Х  | Х  | Х  | Х  | Х  | Х  | Х  |

| Day        | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|------------|----|----|----|----|----|----|----|
| Brigatinib | Х  | Х  | Х  | Χ  | Х  | Х  | Х  |

# About your treatment

Brigatinib blocks proteins that regulate the growth and division of cancer cells, slowing it down.

The tablets are taken once a day continuously. During treatment, your blood tests and any side effects will be monitored.

### How to take the medicines

Swallow the tablets whole with water. Take the tablets at the same time each day with or without a meal. If you miss a dose or vomit, take your usual dose at the next opportunity. Grapefruit or grapefruit juice and St. John's wort must not be taken during treatment.

See also information in the package leaflet.

# Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

## **Fatigue**

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips:Try to exercise and move around. It's the only thing known to help with fatigue.

#### Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

**Tips:**Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

#### Pain in muscles and bones

Pain, tenderness and weakness in the muscles are common.

**Tips:** Can be relieved with common painkillers.

## Numbness and tingling - sensory disturbances

You may experience numbness and tingling in your hands and feet, and your fine motor skills may be impaired. For example, you may find it difficult to press buttons or walk. Symptoms usually go away, but not completely in everyone. Tell your doctor or nurse if you have numbness or tingling in your hands and feet.

#### Skin problems

You may experience itching and a rash.

**Tip**: When washing your skin, be gentle and use mild soap. Lubricate with moisturiser.

## Eye problems

You may get red eyes with stinging and pain. You may also experience blurred vision, dry eyes and become more sensitive to light.

Tip: If your eyes sting and feel dry, use lubricating eye drops available from pharmacies.

## **Hypertension**

High blood pressure is common. Your blood pressure will be monitored during treatment, and you may need medication.

## Contact your clinic immediately at:

Acute

- Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath Contact your reception immediately at:
  - Fever (temp above 38 degrees) or at a temperature below 36 degrees
  - Sudden deterioration regardless of symptom
  - Increased number of stools/diarrhea with or without stomach pain, bloody or black stools
  - New onset dry cough and shortness of breath
  - · Severe itching, blisters, oozing sores and peeling skin.
  - If you or your relatives feel that you are changed, confused or that consciousness is affected.
  - Headache, unusual fatigue, pain behind the eyes and visual disturbances.
  - · Dark foamy urine or very cloudy urine.
  - · Severe muscle or joint pain or an effect on strength or feeling in the arms and legs
  - · Yellow skin color

#### Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe, non-hormonal contraceptives during treatment and for at least 90 days after the end of treatment, as the drug may render the contraceptive pill ineffective.

## General information about cancer

|        |               |          |            |             |         | 4 4 7 7 3 4 |            |          |            |
|--------|---------------|----------|------------|-------------|---------|-------------|------------|----------|------------|
| ΠCΔtii | I information | about ca | ncar ic s  | Mallahla tr | am hath | 11//W       | ardailidan | and Car  | 100rtondor |
| COCILI |               | annin ca | 11651 15 6 | างดแดเมธาน  |         | 11// V      | aiuuuucii  | מווע טמו | 11.121.11  |

www.1177.se

www.cancerfonden.se

#### **Notes**

\_\_\_\_\_\_