Krizotinib

for Lungcancer

Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Krizotinib Morning dose	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Krizotinib Evening dose	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Day	22	23	24	25	26	27	28
Krizotinib Morning dose	X	X	X	X	X	X	X
Krizotinib Evening dose	X	X	X	X	X	X	X

About your treatment

Krizotinib (Xalkori) blocks proteins that regulate the growth and division of cancer cells, slowing it down.

The capsules are taken twice a day (morning and evening) without interruption. During treatment, your blood tests and any side effects will be monitored.

How to take the medicines

Swallow the capsules whole with water. If you miss a dose, take it only if there are more than 6 hours left until your next dose. In case of vomiting, take your usual dose at the next opportunity. You must not eat or drink grapefruit or preparations containing grapefruit or St. John's wort during treatment as this may affect the effect of Krizotinib.

See also information in the package leaflet.

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common in connection with cancer and its treatment. The severity of the symptoms and their duration vary from person to person. You may continue to feel tired long after treatment has ended. This type of extreme exhaustion is known as fatigue—a tiredness that does not always improve with rest or sleep. Many describe fatigue as a total lack of energy, and it can occur even without physical exertion. Your ability to remember, concentrate, and solve problems may be impaired. Fatigue is sometimes associated with feelings of resignation, stress, anxiety, and depression. Tip: Try to stay active. Physical activity often helps alleviate tiredness and fatigue. Find a balance between activity, rest, and sleep that works for you.

Eye problems

You may experience visual disturbances with double vision and flashes of light. You may also get blurred vision and become more sensitive to light. Contact your doctor if you have any symptoms.

Tip: If your eyes sting and feel dry, use lubricating eye drops available from pharmacies.

Swollen legs and feet

Swollen legs and feet are common.

Tips: You can use support stockings. Put your feet up on a footstool or rest for a while in bed. Apply a moisturiser to protect your skin, and watch out for sores.

Problems with stomach and intestines

Constipation is common. You may also get diarrhoea.

Tip: Drink a lot, a few glasses more than usual per day. If you're constipated, try high-fibre foods and exercise. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

Contact your clinic immediately at:

Acute

• Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath

Contact your reception immediately at:

- Fever (temp above 38 degrees) or at a temperature below 36 degrees
- Sudden deterioration regardless of symptom
- Increased number of stools/diarrhea with or without stomach pain, bloody or black stools
- New onset dry cough and shortness of breath
- Severe itching, blisters, oozing sores and peeling skin.
- If you or your relatives feel that you are changed, confused or that consciousness is affected.
- Headache, unusual fatigue, pain behind the eyes and visual disturbances.
- Dark foamy urine or very cloudy urine.
- Severe muscle or joint pain or an effect on strength or feeling in the arms and legs
- Yellow skin color

Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes

_____

_____