

# Tepotinib

for Lungcancer

## Patient information

### Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Tepotinib	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Day	22	23	24	25	26	27	28
Tepotinib	X	X	X	X	X	X	X

## Tepotinib

Tepotinib blocks proteins that regulate the growth and division of cancer cells so that it is slowed down. Tepotinib is taken as tablets. You take the tablets continuously once a day. Your doctor or nurse will tell you how many tablets you should take at a time. During the treatment, your blood samples and any side effects are monitored.

## Tepotinib

Swallow the tablets whole with water. Take the tablets with a meal, at approximately the same time each day. If you miss a dose or vomit, take another dose if it is more than 8 hours until the next scheduled dose. You must not eat or drink preparations with St. John's wort during the ongoing treatment. Read more in the package leaflet.

## Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

### Swollen legs and feet

Swollen legs and feet are common.

**Tips:** You can use support stockings. Put your feet up on a footstool or rest for a while in bed. Apply a moisturiser to protect your skin, and watch out for sores.

### Stomach and intestinal problems

Diarrhea, nausea and vomiting are very common, but you may also become constipated.

### Dry cough and difficulty breathing

New symptoms of dry cough, shortness of breath and difficulty breathing may be due to a special inflammation of the lungs and airways (pneumonitis).

## **Contact your clinic immediately at:**

### **Acute**

Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath

### **Call your clinic immediately at:**

Sudden deterioration regardless of symptoms

Severe diarrhea that does not decrease with stopping drugs

New skin rashes over large parts of the body

Painful skin redness/rash/nail changes

## **Common advice**

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

## **General information about cancer**

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## **Notes**

---

---