# Lenalidomid-Cyklofosfamid-Dexametason

for Haematological malignancy

#### Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Lenalidomid	Х	Х	х	Х	х	х	Х	х	Х	х	Х	Х	Х	Х	Х	Х	х	Х	Х	х	Х
Dexametason	Х							х							Х						
Cyklofosfamidmono- hydrat	х	х	х	Х	х	х	х	х	х	х	Х	х	х	Х	Х	Х	х	Х	х	х	х

#### Next course of treatment, day 29

Day	22	23	24	25	26	27	28
Lenalidomid							
Dexametason	Х						
Cyklofosfamidmono- hydrat							

## About your treatment

Cyclophosphamide is a cytotoxic drug. Lenalidomide nuances your immune system to be able to influence the growth and division of cancer cells. Dexamethasone is a cortisone preparation. Each course is 28 days. Lenalidomide and Cyclophosphamide are taken on days 1-21. Dexamethasone is taken as tablets on days 1, 8, 15 and 22. During the treatment, your blood samples and any side effects are monitored.

#### How to take the medicines

Lenalidimide: Swallow the capsules with water, with or without a meal, at about the same time each day. If you miss a dose, it should only be taken if it is more than 12 hours until the next dose. Cyclophosphamide and Dexamethasone: Swallow the tablets whole with water. Please take the tablets at the same time every day. If you miss a dose or vomit, take your usual dose at the next time. The risk of nausea and vomiting increases when consuming alcohol. You must not eat or drink preparations with St. John's wort during the ongoing treatment. See also information in the package leaflet.

# Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

### **Fatigue**

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips:Try to exercise and move around. It's the only thing known to help with fatigue.

#### Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

**Tip**: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

#### Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

**Tips:**Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

### Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

**Tips:**Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

#### Skin problems

Dry skin and rashes with or without itching are common. Some skin conditions can be made worse by treatment, such as psoriasis and rosacea. You may also get inflammation of the skin (dermatitis), which in rare cases can lead to a severe skin reaction with watery sores.

**Tip for mild itching**: When washing your skin, be gentle and use unscented soap or shower oil. Lubricate with moisturiser.

## Numbness and tingling - sensory disturbances

You may experience numbness and tingling in your hands and feet, and your fine motor skills may become worse. For example, you may have difficulty buttoning buttons or walking. The symptoms usually go away, but not completely in everyone. Tell your doctor or nurse if you have numbness or tingling in your hands and feet. Physical activity can prevent neuropathy and can relieve symptoms and improve motor skills and balance. Try to find the exercise that suits you. Cardio and strength training can be good for preventive purposes. If neuropathy has occurred, improved muscle strength, balance training and yoga can also have an effect by affecting function for the better.

#### Pain in muscles and bones

Pain, tenderness and weakness in the muscles are common.

Tips: Can be relieved with common painkillers.

## Contact your clinic immediately at:

- fever above 38 degrees, or temperature below 36 degrees
- · severe diarrhoea, or diarrhoea combined with fever
- · chest pain or difficulty breathing
- · nosebleeds or other bleeding that you can't stop
- · skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- · chest pain or difficulty breathing, call 112

## Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden: www.1177.se

www.cancerfonden.se

Notes	
-------	--