

Regimen schedule

Day	Next course of treatment, day 22																				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Ifosfamid	X	X																			
Mesna	X	X																			
Vinkristin	X																				
Daktinomycin	X																				
Pegfilgrastim			X																		

About your treatment

Ifosfamide, vincristine and dactinomycin are cytostatics. Mesna is given to protect the urinary tract. Also included is filgrastim, which stimulates the bone marrow to produce white blood cells and strengthens your immune system. Each course is 21 days. Ifosfamide is given as a drip on days 1 and 2. Vincristine and dactinomycin are given as a drip on day 1. Mesna is given via a syringe on days 1 and 2. Filgrastim is given via a syringe on days 5-12. Sometimes you can take the injections yourself. Alternatively, a special form of filgrastim is used, which is then given on only one occasion. During the treatment, your blood samples and any side effects are monitored

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips: Try to exercise and move around. It's the only thing known to help with fatigue.

Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

Tip: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

Tips: Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

Numbness and tingling - sensory disturbances

You may experience numbness and tingling in your hands and feet, and your fine motor skills may become worse. For example, you may have difficulty buttoning buttons or walking. The symptoms usually go away, but not completely in everyone. Tell your doctor or nurse if you have numbness or tingling in your hands and feet. Physical activity can prevent neuropathy and can relieve symptoms and improve motor skills and balance. Try to find the exercise that suits you. Cardio and strength training can be good for preventive purposes. If neuropathy has occurred, improved muscle strength, balance training and yoga can also have an effect by affecting function for the better.

Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

Tips: Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

Bleeding

You may have an increased risk of bleeding. For example, you may bleed easily from your nose or mouth. If you get a cut on your skin, you may bleed longer than usual. Surgical wounds may heal more slowly.

Tips: Be careful not to cut yourself. You can stop nosebleeds by squeezing the soft parts of your nose for 10 minutes. If this doesn't stop it, contact your doctor. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist or other health care provider of your cancer treatment when you visit.

Problems with skin, hair and nails

Skin often becomes dry and more sensitive to the sun. You may also lose hair or your hair may change texture. Nails can become brittle.

Tip: When washing your skin, be gentle and use mild soap. Lubricate often with moisturiser. Wear covering clothing to protect against the sun, and sunscreen on parts of your body that you can't cover. Wear gloves for work that is dirty or abrasive for your hands. If your nails are bothering you, you can use nail polish and brush your cuticles with oil or apply a moisturiser.

Problems with mouth and mucous membranes

The mucous membranes of your nose, mouth, eyes and abdomen may be affected by your treatment. For example, you may have nosebleeds and a runny nose. In the mouth, you may experience dryness, redness, burning and sores. You can also get fungus, which turns the lining of your mouth red or gives it a white coating. Eyes may become dry and watery. The mucous membranes in the lower abdomen can also become dry.

Tip: If you have a dry mouth, use saliva stimulants available in pharmacies. Rinsing your mouth with Vichy water can prevent fungus. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist of your cancer treatment when you visit. Avoid contact lenses if you have eye problems. Women can use oestrogen cream, emollient cream or oil to treat dry mucous membranes in the lower abdomen.

Pain in muscles and bones

Pain, tenderness and weakness in the muscles are common.

Tips: Can be relieved with common painkillers.

Contact your clinic immediately at:

- fever above 38 degrees, or temperature below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- chest pain or difficulty breathing
- nosebleeds or other bleeding that you can't stop
- skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

Common advice

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes
