

**Avelumab-Axitinib**

for Urological cancer

**Regimen schedule**

Next course of treatment, day 15

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Avelumab	X													
Axitinib Morning dose	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Axitinib Evening dose	X	X	X	X	X	X	X	X	X	X	X	X	X	X

**About your treatment**

Avelumab is an antibody that inhibits the growth of cancer cells. Axitinib blocks proteins that regulate the growth and division of cancer cells, slowing them down.

Each course is 14 days. Day 1, you get an IV. Axitinib (Inlyta) is taken as tablets twice a day on days 1-14.

During treatment, your blood tests and any side effects will be monitored.

**Side effects - symptoms and tips**

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. The side effects may require a break in treatment. Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

**Side effects may come late, several months after treatment has ended. Contact your doctor anyway!**

**Side effects - symptoms and tips**

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

**Fatigue**

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

**Tips:** Try to exercise and move around. It's the only thing known to help with fatigue.

**Nausea and taste changes**

You may feel unwell from the treatment. You will therefore be given anti-nausea medication in conjunction with the treatment. The medicines may cause constipation. You can get different varieties depending on how you feel. The taste of food and drink may be altered by the treatment.

**Tips:** Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

## **Problems with stomach and intestines**

Diarrhoea is common. You may also become constipated.

**Tip:** Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

## **Hypertension**

High blood pressure is common. Your blood pressure will be monitored during treatment, and you may need medication.

## **Skin and nails**

Skin often becomes dry and more sensitive to the sun. Nails can become brittle.

**Tips:** When washing your skin, be gentle and use mild soap. Lubricate with moisturiser. Wear covering clothing to protect against the sun, and sunscreen on parts of your body that you can't cover. Wear gloves for work that is dirty or abrasive for your hands. If your nails are bothering you, you can use nail polish and brush your cuticles with oil or apply a moisturiser.

## **Problems with hands and feet**

You may sometimes experience redness, dry flaky skin and swelling on the soles of your hands and feet. It usually starts as soreness, tingling and numbness, which can develop into blisters, cracks and sores. Contact your doctor immediately if you experience these symptoms.

**Tip:** Avoid exposing hands and feet to high heat, such as showering or washing dishes with very hot water. To relieve the discomfort, you can use cold items, such as a freezing pack wrapped in a towel. Also avoid exposing your hands and feet to vibration or abrasion, such as when using a drill for hours or running for miles.

## **Headache**

Headaches are common.

## **Bleeding**

You may have an increased risk of bleeding. For example, you may bleed easily from your nose or mouth. If you get a cut on your skin, you may bleed longer than usual. Surgical wounds may heal more slowly.

**Tips:** Be careful not to cut yourself. You can stop nosebleeds by squeezing the soft parts of your nose for 10 minutes. If this doesn't stop it, contact your doctor. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist or other health care provider of your cancer treatment when you visit.

## **Cough and shortness of breath**

You may experience coughing and shortness of breath. Tell your doctor or nurse if you have trouble breathing.

## **Pain in muscles and joints**

Pain in muscles and joints is common.

## **Contact your clinic immediately at:**

- fever over 38 degrees or if your temperature drops below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

## Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## Notes

---

---