

**Cetuximab-FOLF**

for Gastrointestinal cancer

**Regimen schedule**

Day	Next course of treatment, day 15													
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Cetuximab	X													
Kalciumfolinat (vattenfritt)	X													
Fluorouracil	X													
Fluorouracil Bärbar infusionspump (46 tim.)	→	→	*											

\* Pumpen kopplas bort.

**About your treatment**

Cetuximab is an antibody that inhibits the growth of cancer cells. FOLF is an abbreviation for fluorouracil, which is a cytostatic and calcium folinate, a folic acid that increases the effect of fluorouracil. Each course is 14 days. On day 1, you will receive a drip and a portable infusion pump that doses fluorouracil for 2 days at home. After that, there is a break of 12 days before a new course starts. During the treatment, your blood tests and any side effects are monitored

**Portable infusion pump**

Homepump or Intermate are portable, single-use infusion pumps. The pump is driven by the pressure created when it is filled with a solution. The pump is usually stored in a waist bag. Check that the hose is not pinched. Avoid getting the pump wet. If the pump falls to the floor, it should be able to withstand this. It may be a good idea to check the pump daily and check if it is not free.

**Side effects - symptoms and tips**

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

**Fatigue**

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

**Tips:** Try to exercise and move around. It's the only thing known to help with fatigue.

**Nausea and taste changes**

You may feel unwell from the treatment. You will therefore be given anti-nausea medication in conjunction with the treatment. The medicines may cause constipation. You can get different varieties depending on how you feel. The taste of food and drink may be altered by the treatment.

**Tips:** Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

## **Problems with stomach and intestines**

Diarrhoea is common. You may also become constipated.

**Tip:** Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

## **Acne-like skin problems**

You may get acne-like rashes on your face and upper body. You may also experience itching, cracking of the skin and inflammation around the nails. Talk to your nurse if you develop skin problems - you may need medication.

**Tip:** Apply a moisturising cream to your whole body several times a day. Use bath oil instead of soap. Avoid sun-bathing. Wear a hat and full-coverage clothing. Lubricate parts of the body not covered by clothing with sunscreen that has a high SPF and UVA protection.

## **Problems with mouth and mucous membranes**

The mucous membranes of your nose, mouth, eyes and abdomen may be affected by your treatment. For example, you may have nosebleeds and a runny nose. In the mouth, you may experience dryness, redness, burning and sores. You can also get fungus, which turns the lining of your mouth red or gives it a white coating. Eyes may become dry and watery. The mucous membranes in the lower abdomen can also become dry.

**Tip:** If you have a dry mouth, use saliva stimulants available in pharmacies. Rinsing your mouth with Vichy water can prevent fungus. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist of your cancer treatment when you visit. Avoid contact lenses if you have eye problems. Women can use oestrogen cream, emollient cream or oil to treat dry mucous membranes in the lower abdomen.

## **Eye problems**

You may get red eyes with stinging and pain. You may also experience blurred vision, dry eyes and become more sensitive to light.

**Tip:** If your eyes sting and feel dry, use lubricating eye drops available from pharmacies.

## **Problems with the heart**

The treatment carries a certain risk of cardiac effects. You may have symptoms such as chest pain, an unusual heart rhythm, palpitations, shortness of breath or swollen legs. ECG and/or ultrasound of the heart are sometimes included in checks on heart function.

## **Infection susceptibility**

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

**Tips:** Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

## **Problems with skin, hair and nails**

Skin often becomes dry and more sensitive to the sun. You may also lose hair or your hair may change texture. Nails can become brittle.

**Tip:** When washing your skin, be gentle and use mild soap. Lubricate often with moisturiser. Wear covering clothing to protect against the sun, and sunscreen on parts of your body that you can't cover. Wear gloves for work that is dirty or abrasive for your hands. If your nails are bothering you, you can use nail polish and brush your cuticles with oil or apply a moisturiser.

## Contact your clinic immediately at:

- fever above 38 degrees, or temperature below 36 degrees
- severe diarrhoea, or diarrhoea combined with fever
- chest pain or difficulty breathing
- nosebleeds or other bleeding that you can't stop
- skin rash on most of the body, or skin rash with pain
- sudden deterioration, regardless of symptoms
- chest pain or difficulty breathing, call 112

## Common advice

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

## General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

[www.1177.se](http://www.1177.se)

[www.cancerfonden.se](http://www.cancerfonden.se)

## Notes

---

---