

Niraparib tableter

for Gynecological cancer

Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Niraparib	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Next course of treatment, day 29

Day	22	23	24	25	26	27	28
Niraparib	X	X	X	X	X	X	X

About your treatment

Niraparib (Zejula) blocks proteins that regulate the growth and division of cancer cells, slowing them down.

The capsules are taken once a day without interruption.

During treatment, your blood tests and any side effects will be monitored.

How to take the medicines

Swallow the tablets whole with water once a day, without food (at least 1 hour before or 2 hours after a meal) or with a light meal. Take the tablets at approximately the same time each day, preferably at bedtime. If you miss a dose or vomit, take your usual dose at the next opportunity.

Read more in the package leaflet.

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips: Try to exercise and move around. It's the only thing known to help with fatigue.

Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

Tips: Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

Tip: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

Hypertension

High blood pressure is common. Your blood pressure will be monitored during treatment, and you may need medication.

Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

Tips: Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

Skin problems

You may experience itching and a rash.

Tip: When washing your skin, be gentle and use mild soap. Lubricate with moisturiser.

Pain in muscles and joints

Pain in muscles and joints is common.

Contact your clinic immediately at:

Acute

- Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath

Contact your reception immediately at:

- Fever (temp above 38 degrees) or at a temperature below 36 degrees
- Sudden deterioration regardless of symptom
- Increased number of stools/diarrhea with or without stomach pain, bloody or black stools
- New onset dry cough and shortness of breath
- Severe itching, blisters, oozing sores and peeling skin.
- If you or your relatives feel that you are changed, confused or that consciousness is affected.
- Headache, unusual fatigue, pain behind the eyes and visual disturbances.
- Dark foamy urine or very cloudy urine.
- Severe muscle or joint pain or an effect on strength or feeling in the arms and legs
- Yellow skin color

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes
